


LINEAR ALGEBRA

WEEK 2

Roadmap



FEEDBACK

- a) Let $\mathbf{0} \in \mathbb{R}^m$ denote the vector whose entries are all zero. We say that a set L is a line through the origin in $\mathbb{R}^m$ if and only if there exists $\mathbf{w} \in \mathbb{R}^m$ with $\mathbf{w} \neq \mathbf{0}$ such that $L = \{\lambda \mathbf{w} : \lambda \in \mathbb{R}\}$. Let now L be a line through the origin in $\mathbb{R}^m$ and let $\mathbf{u}$ be an arbitrary non-zero element of L . Prove that $L = \{\lambda \mathbf{u} : \lambda \in \mathbb{R}\}$.

$$\begin{aligned} & \exists \mathbf{u} \in \mathbb{R}^m : L = \{\lambda \mathbf{u} : \lambda \in \mathbb{R}\} \quad \textcircled{1} \\ & \Rightarrow \mathbf{u} = \lambda_{\mathbf{u}} \mathbf{w} \quad \textcircled{2} \end{aligned}$$

$$\text{z.z. } L = \{\lambda \mathbf{u} : \lambda \in \mathbb{R}\}$$

$$\begin{aligned} \text{"} \subseteq \text{" : betrachte } v \in L : v &= \lambda_v \mathbf{w} \\ &\Rightarrow v = \frac{\lambda_v}{\lambda_{\mathbf{u}}} \mathbf{u} \quad \textcircled{2} \\ &\Rightarrow v \in \{\lambda \mathbf{u} : \lambda \in \mathbb{R}\} \end{aligned}$$

$$\begin{aligned} \text{"} \supseteq \text{" : betrachte } v' \in \{\lambda \mathbf{u} : \lambda \in \mathbb{R}\} : v' &= \lambda_{v'} \mathbf{u} \\ &\Rightarrow v' = \lambda_{v'} \mathbf{u} = \lambda_{v'} \lambda_{\mathbf{u}} \mathbf{w} \\ &\Rightarrow v' \in L \end{aligned}$$

- b) For two lines through the origin L_1 and L_2 in $\mathbb{R}^m$, prove that we have either $L_1 \cap L_2 = \{0\}$ or $L_1 \cap L_2 = L_1 = L_2$.

$$L_1 = \{\lambda u_1 : \lambda \in \mathbb{R}\} \textcircled{1}$$

$$L_2 = \{\lambda u_2 : \lambda \in \mathbb{R}\} \textcircled{2} \quad u_1, u_2 \in \mathbb{R}^m$$

$$\text{z.z.: } L_1 \cap L_2 \stackrel{\textcircled{1}}{=} \{0\} \text{ oder } L_1 \cap L_2 \stackrel{\textcircled{2}}{=} L_1 = L_2$$

1) Beide Fälle sind unmöglich, weil $\{0\} \neq L_1$ oder $\{0\} \neq L_2$

$$2) 0 = 0 \cdot u_1 \text{ und } 0 = 0 \cdot u_2 \Rightarrow 0 \in L_1 \text{ und } 0 \in L_2 \\ \Rightarrow 0 \in L_1 \cap L_2$$

3) Wir haben 2 Fälle:

Fall 1 ($L_1 \cap L_2 = \{0\}$):

dies impliziert automatisch $\textcircled{1}$

Fall 2 ($L_1 \cap L_2 \neq \{0\}$):

$$L_1 \cap L_2 \neq \emptyset \Rightarrow \exists u \in \mathbb{R}^m : u \in L_1 \cap L_2 \quad (u \neq 0, \text{ weil } L_1 \cap L_2 \neq \{0\})$$

$$\left. \begin{array}{l} u \in L_1 \stackrel{\textcircled{1}}{\Rightarrow} L_1 = \{\lambda u : \lambda \in \mathbb{R}\} \\ u \in L_2 \stackrel{\textcircled{2}}{\Rightarrow} L_2 = \{\lambda u : \lambda \in \mathbb{R}\} \end{array} \right\} \Rightarrow L_1 = L_2 = L_1 \cap L_2 \textcircled{2}$$

Da wir alle Fälle abgedeckt haben und entweder $\textcircled{1}$ oder $\textcircled{2}$ gilt, gilt die Aussage als bewiesen.

c) Consider an arbitrary line through the origin L in $\mathbb{R}^2$. Prove that L is a hyperplane, i.e. find a vector $d \neq 0$ such that

$$L = \{v \in \mathbb{R}^2 : v \cdot d = 0\}.$$

↳ per Def.: $\exists u = \begin{pmatrix} u_1 \\ u_2 \end{pmatrix} : L = \{\lambda u : \lambda \in \mathbb{R}\}$ ①
↳ wir suchen $d = \begin{pmatrix} d_1 \\ d_2 \end{pmatrix}$, sodass $u \cdot d = 0$
 $\Rightarrow d_1 := -u_2; d_2 := u_1$ ②

$$\text{z.z.: } L = \{v \in \mathbb{R}^2 : v \cdot d = 0\}$$

$$"\subseteq": \text{ wähle } u \in L : u = \lambda u \quad ①$$

$$u \cdot d = (\lambda u) \cdot d = \lambda(u \cdot d) \stackrel{②}{=} \lambda \cdot 0 = 0$$

$$\Rightarrow u \in \{v \in \mathbb{R}^2 : v \cdot d = 0\}$$

$$"\supseteq": \text{ wähle } u \in \{v \in \mathbb{R}^2 : v \cdot d = 0\}$$

$$u \cdot d = u_1 d_1 + u_2 d_2 = 0$$

$$\stackrel{②}{\Rightarrow} u \cdot d = -u_1 u_2 + u_2 u_1 = 0$$

$$\Rightarrow u_2 u_1 = u_1 u_2$$

$$\Rightarrow u_2 = \frac{u_2}{u_1} u_1 = \frac{u_2}{u_1} u_2$$

für $u_1 \neq 0$:

$$\text{wähle } \lambda = \frac{u_2}{u_1} \Rightarrow u_1 = \lambda u_1 \text{ und } u_2 = \lambda u_2$$

$$\Rightarrow u = \lambda u$$

$$\Rightarrow u \in L \quad ①$$

für $u_2 \neq 0$: Beweis analog

↳ wir müssen entweder $u_1 \neq 0$ oder $u_2 \neq 0$ Fälle prüfen
wegen $u \neq 0$ ③

↑ Proof by case distinction (mehr im DiskMath Skript in Kapitel "Proof Patterns")

FRAGEN?

Definition 2.10 ((In)dependent column and (column) rank of a matrix). *Let*

$$A = \begin{bmatrix} | & | & \cdots & | \\ \mathbf{v}_1 & \mathbf{v}_2 & \cdots & \mathbf{v}_n \\ | & | & \cdots & | \end{bmatrix}$$

be an $m \times n$ matrix with columns $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n \in \mathbb{R}^m$. Column $\mathbf{v}_j$ is called independent if $\mathbf{v}_j$ is not a linear combination of $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_{j-1}$. Otherwise, $\mathbf{v}_j$ is called dependent. The rank of A , written as $\text{rank}(A)$, is the number of independent columns of A . The rank is sometimes also called the column rank to distinguish it more explicitly from the row rank, see Definition 2.14 below.

In-class

1. Rank of a matrix (in-class) (★★☆)

Let $m \in \mathbb{N}_{\geq 2}$ be arbitrary and consider the $m \times m$ matrix

$$A_m = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1m} \\ a_{21} & a_{22} & \dots & a_{2m} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mm} \end{bmatrix}$$

with $a_{ij} = i + j$ for all $i, j \in \{1, 2, \dots, m\}$.

a) Calculate A_m for $m \in \{2, 3, 4\}$.

In-class

$$A_2 = \begin{bmatrix} 2 & 3 \\ 3 & 4 \end{bmatrix}, A_3 = \begin{bmatrix} 2 & 3 & 4 \\ 3 & 4 & 5 \\ 4 & 5 & 6 \end{bmatrix}, \text{ and } A_4 = \begin{bmatrix} 2 & 3 & 4 & 5 \\ 3 & 4 & 5 & 6 \\ 4 & 5 & 6 & 7 \\ 5 & 6 & 7 & 8 \end{bmatrix}.$$

In-class

1. Rank of a matrix (in-class) (★★☆)

Let $m \in \mathbb{N}_{\geq 2}$ be arbitrary and consider the $m \times m$ matrix

$$A_m = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1m} \\ a_{21} & a_{22} & \dots & a_{2m} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mm} \end{bmatrix}$$

with $a_{ij} = i + j$ for all $i, j \in \{1, 2, \dots, m\}$.

- a) Calculate A_m for $m \in \{2, 3, 4\}$.
- b) Determine the rank of A_m . You need to motivate your answer.

Aufgaben

a) Sei $A \in \mathbb{R}^{m \times n}$ mit $A = [\overset{\textcircled{1}}{a_1} \dots a_n]$
zeige : $0 \in C(A)$ und $\forall j \in [n] : a_j \in C(A)$ $\textcircled{2}$

Aufgaben

b) Sei $A \in \mathbb{R}^{m \times n}$

zeige:

① $\forall y, z \in C(A) \forall \lambda \in \mathbb{R} : y + \lambda z \in C(A)$

② $\forall u, v \in N(A) \forall \lambda \in \mathbb{R} : u + \lambda v \in N(A)$

Aufgaben

c) Sei $A \in \mathbb{R}^{m \times n}$ und $b \in \mathbb{R}^m$. Es gilt für $x_0 \in \mathbb{R}^n$ $Ax_0 = b$
zeige: $\{x \in \mathbb{R}^n : Ax = b\} = \{x_0 + z : z \in N(A)\}$

Proof: $(Ax)^T = x^T A^T$, $A \in \mathbb{R}^{m \times n}$ $x \in \mathbb{R}^n \Rightarrow Ax \in \mathbb{R}^m$

$\swarrow$ oder $\mathbb{R}^{n \times 1}$ $\swarrow$ oder $\mathbb{R}^{m \times 1}$

① $(Ax)_{i1} = \sum_{j=1}^n a_{ij} x_j \Rightarrow (Ax)^T_{1i} = \sum_{j=1}^n a_{ij} x_j$

② $(A^T)_{ji} = a_{ij} \Rightarrow (x^T A^T)_{1i} = \sum_{j=1}^n x_j (A^T)_{ji} = \sum_{j=1}^n x_j a_{ij}$

$\boxed{0} = \text{---} \boxed{0}$

$(Ax)^T_{1i} = \sum_{j=1}^n a_{ij} x_j = \sum_{j=1}^n x_j a_{ij} = (x^T A^T)_{1i}$
 $\Rightarrow (Ax)^T = x^T A^T$

KAHOOT